## Observing Gravitational Waves from the Next Nearby Core-Collapse Supernova


Sarah Gossan<sup>[1]</sup>, Christian Ott<sup>[1]</sup>, Ernazar Abdikamalov<sup>[1]</sup>, Alexandra DeMaio<sup>[2,1]</sup>, Erik Katsavounidis<sup>[3]</sup>

[1] California Institute of Technology, [2] Rutgers, [3] Massachusetts Institute of Technology


TAUP 2013, 12th September 2013

## Prospects for Nearby Core-Collapse Supernovae

- \* Important galaxies for CCSN rate in Local Group (CCSNe/100yr)
  - \* Milky Way (0.5-2.5)
  - \* LMC (0.1-0.5)
  - \* NGC6822 (~0.04)
  - \* M31 (0.2-1.2)
- \* M82 @ 3.52Mpc with (3.0-13.0) CCSNe/100yr
- CCSNe most likely to be ~10kpc from galactic center.




#### The Advanced Detector Network



## Challenges in Detecting CCSNe

- \* Low event rate for aLIGO.
- \* Duty cycle not optimal.
- \* Non-Gaussian and nonstationary detector noise.
- \* Can't predict exact GW signal expected robustly.




Detector	S5y1 (Nov 05-Nov 06) Duty Cycle
H1	72.8%
L1	59.3%


Abbott+09, Phys. Rev. D80:102001

#### The Excess-Power Search


- \* Uses time-frequency signal content.
- \* Optimal for unmodelled signals.
- \* Does data contains excesspower than expected from noise?
- \* Increase accuracy by reducing `on-source' region (OSR).
- \* X-Pipeline (Sutton+09) and Coherent WaveBurst (Klimenko+08).




## Utilizing Multi-Messenger Observations




#### GW Searches for CCSNe so far



#### A Joint GW- $\nu$ Search for CCSNe



### Detecting CCSNe with aLIGO



- \* Unlikely to detect average CCSNe beyond Milky Way/LMC/SMC.
- Non-detections
  make statements
  about
  progenitors.
- Sub-threshold signals in GW and neutrinos combine to increase observational evidence.


## Utilizing sub-threshold GW/ $\nu$ signals

\* GRB051103.

\* Estimate noise background.

\* Exclude GW emission models.


\* Improve neutrino search sensitivity - require GW coincidence.



#### Parameter Estimation for CCSNe

\* Want astrophysically interesting information from observations.

\* Don't have exact analytical expressions - Principal Component Analysis (PCA) captures dominant signal features.





- \* Reconstruct low SNR signals.
  - Röver+09.

\* Determine CCSN explosion mechanism. Logue+12.

### PCA and Bayesian Inference

- \* PCA create basis sets encompassing `principal components' of simulation waveform space.
- \* Generate PCs from time-domain, linearly polarized waveforms.
- \* Bayesian Inference with Nested Sampling algorithm.
  - \* PE: Estimate PC coefficients and reconstruct injected signals using PCs..
  - \* Model Selection: calculate evidence for each model which PC set is most ----- likely given observed data?







# Inferring the Differential Rotation of CCSN Progenitors

Ernazar Abdikamalov, Sarah Gossan, Alexandra DeMaio, Christian Ott


- \* Characterize progenitor rotation for rotating corecollapse sources.
- \* Optimal-orientation and source at 10kpc Only A and  $\beta_{ic,b}$  unknown.
- \* Two approaches
  - \* Model selection.
  - \* Numerical template bank.



#### Determining A with Model Selection



# A Numerical Template Bank - Matched Filtering



#### Future Work

- \* Prepare for the first detection of GW from CCSNe!
  - \* More collaboration between GW and  $\nu$  working groups.
  - \* Low-latency burst pipelines.
  - \* Real-time signal analysis.
  - \* Extend differential rotation analysis to additional physical progenitor parameters, e.g. different explosion mechanisms, equation of state.
- \* Inference analysis for both GW and  $\nu$  signatures of CCSNe.